

ROLE OF POST-OP TOPICAL DILTIAZEM CREAM IN PAIN CONTROL AFTER MILLIGAN-MORGAN HEMORRHOIDECTOMY

Rashid Aslam, Zeeshan Saboor Ahmad, Nayab Iqbal, Tasmia, Aneela Khan, Muhammad Fawad Khan, Fida Hussain

ABSTRACT

Background: Spasm of internal anal sphincter is thought to be responsible for pain after hemorrhoidectomy. Topical use of a calcium channel blocker, 2% Diltiazem (DTZ), may be effective in reducing pain after hemorrhoidectomy.

Objective: To evaluate and compare pain scores on Visual Analogue Scale post-operatively after hemorrhoidectomy from day two to day five, in patients using 2% DTZ versus patients using Placebo ointment along with standard treatment for pain relief. Need for urinary catheterization was also assessed indicating severity of pain causing urinary retention.

Materials and Methods: This was a randomized control trial conducted at Hayatabad Medical Complex, Peshawar, for a period of fourteen months from 01st June 2016 to 31st August, 2017. One hundred patients, who had undergone Milligan Morgan hemorrhoidectomy, were randomly assigned to receive 2% DTZ ointment (Group A - study Group) or a placebo ointment-petroleum jelly (Group B - control group) post-operatively. Fifty patients were included in each group. Postoperative pain scores were recorded using Visual Analogue Score on second postoperative day. SPSS version 16 was used for statistical analysis.

Results: Mean age was 35.1 ± 13.3 years. 64(64%) patients were males whereas thirty six (36%) were females, with male to female ratio being 1.7:1. Patients using 2% DTZ ointment had significantly less pain after surgery. The best pain relief in range of 1-3 on Visual Analogue Scale was achieved in 36% patients in group-A(Study Group) while the same score was achieved by only 12% of patients in group-B(Control Group). Urinary retention because of severe pain occurred only in 2 patients(0.4%) in group-A while it occurred in 6 patients(12%) in group-B, for which catheterization was done.

Conclusion: Perianal application of 2% Diltiazem ointment after hemorrhoidectomy is effective in reducing pain post-operatively.

Key Words: Hemorrhoidectomy, Calcium Channel Blockers, Diltiazem, Milligan Morgan, hemorrhoidectomy

INTRODUCTION

Haemorrhoids are the pathological enlargement and distal displacement of the normal anal cushions and dysregulation of the vascular tone and vascular hyperplasia seems playing important role in haemorrhoidal development and could be a potential target for medical treatment¹.

Hemorrhoidectomy is the only truly curative treatment to reduce recurrent symptoms in patients with grade 3 or 4 haemorrhoids². Indications for hemorrhoidectomy include failure of non operative management, acute complicated haemorrhoids such as strangulation or thrombosis, patient preference and concomitant anorectal conditions³.

Milligan Morgan Hemorrhoidectomy⁴ is considered a relatively safe surgical treatment for the disease but complications exist. Pain at the operative site is a

Department of Surgery MTI, HMC Peshawar - Pakistan

distressing complication after this procedure resulting in anxiety, constipation, urinary retention and increased hospital stay^{5,6}.

Warm sitz bath, stool softeners and systemic analgesics are standard post-operative regimen to reduce post-operative pain⁷. Spasm of internal anal sphincter is considered as one of the main factor contributing towards pain^{4,8}. Internal sphincterotomy along with hemorrhoidectomy has been employed to reduce spasm but it has been found to be associated with faecal incontinence (upto 8-30%)^{9,10}.

Internal anal sphincter has a calcium dependent mechanism to maintain its tone. Diltiazem (DTZ) a calcium channel blocker, blocks calcium uptake by the myocytes, thus relaxing the smooth muscles of internal anal sphincter⁵. Due to fewer side effects, 2% DTZ is replacing topical glyceryl trinitrate in treatment of chronic anal fissure¹¹.

The rationale behind this study was the conflict of data and lack of scientific consensus on selecting the best modality for post-op pain control after Milligan-Morgan hemorrhoidectomy.

Address for correspondence:

Dr. Rashid Aslam

Department of Surgery MTI, HMC Peshawar - Pakistan
Cell: 0333-3944949

MATERIALS AND METHODS

This study was a randomized controlled trial conducted at Department of General Surgery, Hayatabad Medical Complex Peshawar.

All patients with third and fourth degree haemorrhoids admitted to the ward from 1st June 2016 to 31st August 2017 were included after evaluation by consultant, informed consent was taken from each patient about surgery and this study. Using non-probability purposive sampling (by random allocation employing flip coin method), 100 patients with 3rd and 4th degree haemorrhoids were included in the trial and divided equally into two groups i.e. Group-A allocated to treatment by topical 2% DTZ (study group) and Group-B topical placebo (control group).

Inclusion Criteria

- i. Patients having 3rd and 4th degree haemorrhoids
- ii. Aged \geq 16 years and \leq 70 years

Exclusion Criteria

- i. Previous hx of anal procedure/ pathology, anal fissure, fistula in ano, thrombosed hemorrhoids and malignancy.
- ii. Patients in whom calcium channel blocker are contraindicated or any other allergy or co-morbidity.
- iii. Patients with diabetes mellitus immunosuppression or other spinal/ neurological abnormalities.

Standard post-operative management was employed in patients of both groups. These included IV Metronidazole 40mg TDS for three days; analgesia in the form of Ketorolac IV 30 mg TDS for three days. Stool softness syrup Cremafin 2 TSF in TDS dose daily were given. Warm sitz bath for 15 minutes twice daily after immediate post-operative day was taken by all the patients.

Anal packing was removed after 24 hours. In Group-A, patients were advised to apply 2% DTZ ointment via applicator to the perianal region twice daily for four days; first dose being applied on first post-op day after removal of pack. In Group-B, placebo ointment (petroleum jelly) was advised to be applied accordingly. Both ointments looked same and were supplied in dispensers of identical appearance. All patients were examined by the consultant who was unaware of the study group of the patients on every morning of post-operative day; from day 2 to 5.

Post-operative pain was measured by Visual Analogue Score and need of catheterization because of severe pain.

Self-designed questionnaire was used to document all findings. It included demographic data of the patients and average of pain scores on Visual Analogue

Scale and the need of catheterization.

The software program SPSS was utilized for all statistical analysis. The p-value of 0.05 was considered significant statistically.

RESULTS

The age range of patients was between 16 to 17 years with overall mean (\pm SD) age of patients 35.1 (\pm 13.3) years. Out of 100 patients in this study, 64 (64%) were male and 36 (36%) were female; and male to female ratio was 1.7:1 in figure 1. Mean pain scores (percentage) as assessed on Visual Analogue Score and the need of catheterization because of urinary retention called caused by severe pain.

As evident from results, best pain relief (1-3 on Visual Analogue Scale) was achieved in 18 (36%) of patients in study Group-A while only in 6 (12%) patients in control Group-B. The same is the case with urinary catheterization which was required by only 2 (0.04%) patients in study Group-A while needed by 6 (12%) patients in control Group-B because of urinary retention caused by severe pain.




Figure 1

Table 1

Pain Score	Group-A n(%)	Group-B n(%)
1-3	18 (0.36)	6 (0.12)
4-5	20 (0.40)	26 (0.52)
6-8	9 (0.18)	12 (0.24)
9-10	3 (0.06)	6 (0.12)
Catheterization	2 (0.04)	6 (0.12)

DISCUSSION

Post-hemorrhoidectomy pain has always been a major concern both for the surgeons and patients¹². It is associated with hypertonia of internal sphincter¹³ and the lateral sphincterotomy has been found effective in muscle relaxation, pain relief and quick healing¹⁴. Galzia et al¹⁵ prospectively randomized 42 consecutive patients with prolapsed haemorrhoids concluded that lateral sphincterotomy with hemorrhoidectomy significantly improves pain as compared to hemorrhoidectomy alone. Kanellos and associates⁹ evaluated the effect of lateral sphincterotomy on pain after Milligan-Morgen hemorrhoidectomy and concluded that the addition of lateral sphincterotomy to open hemorrhoidectomy have a positive effect as it relaxes muscles but may be associated with significant risk of incontinence¹⁰.

Topical application of nitroglycerine has also been employed by various surgeons¹⁶ after disappointment of long term result of lateral sphincterotomy^{10,6}. The pain post-operatively was reduced¹⁶ but frequency of headache was high and required additional analgesics. With understanding of intestinal and sphincter physiology, alternative treatment with fewer side effects is necessitated.

Diltiazem is a CCB with few side effects and similar benefits as compared to nitroglycerine¹⁷⁻¹⁹. Amoli et al conducted a prospective study to evaluate the role of topical 2% Diltiazem after Hemorrhoidectomy and came to the conclusion that topical Diltiazem cream is of significant importance in reducing post-operative pain²⁰.

Similar study was conducted by Silvermen et al⁵ using DTZ post-Hemorrhoidectomy for pain relief and they concluded that topical DTZ reduces pain and other co-morbidity. Chauhan et al prospectively randomized 108 patients of Grade 3 & 4 haemorrhoids and compared the efficacy of internal sphincterotomy with topical application of 2% DTZ ointment after Hemorrhoidectomy for the relief of pain. No significant difference was found in mean pain scores as both procedures involved relieving the spasm of anal sphincter while the former one is more radical.

In our study, pain perception was significantly reduced in the Diltiazem group as compared to Placebo group. Best pain relief (1-3 on Visual Analogue Scale) was achieved in 18 (36%) of patients in study Group-A while the same level of pain relief was achieved in Placebo Group-B in only 6 patients (12%).

The urinary catheterization was only needed in two patients out of fifty patients in study Group-A while six patients out of fifty patients needed urinary catheterization in Placebo Group-B (control group) which showed the severity of pain causing urinary retention.

CONCLUSION

It is concluded that application of 2% Diltiazem ointment along with standard treatment significantly reduces the pain after Hemorrhoidectomy (Morgan-Milligan Hemorrhoidectomy) as compared to other radical procedures. On the basis of above study, topical DTZ 2% ointment can be used along with standard post-op regimen after Hemorrhoidectomy; but it still needs a more robust study and comparison with other modalities.

REFERENCES

1. Lohsiriwat, V., 2012. hemorrhoids from basic pathophysiology to clinical management. *World J. Gastroenterol.*, 18 : 2009 – 2017.
2. Mounsey, A.L., J. Halladay and T.S . Sadiq, 2011. Hemorrhoids. *Am Family Physician*, 84 : 204 – 210.
3. Mirfazaelian, H, S. Nikfar, S. Derakhshani and M. Abdollahi, 2012. Efficacy and incontinence rate of biomaterials (fibrin glue and fibrin plug) in comparison to surgical interventions in the treatment of perianal fistula : A systematic review and meta-analysis of randomised controlled trials. *Int. J. Pharmacol.*, DOI : 10.3923/ijp.2012.
4. Hunt L, Luck A J, Rudkin G, Hewet PJ. Day- case hemorrhoidectomy. *Br J Surg* 1999; 86: 255 – 8.
5. Silverman R, Bendick PJ, Wasvarg H J. A randomised, prospective, double blind, placebo-controlled trial of the effects of a calcium channel blocker ointment on pain after hemorrhoidectomy. *Dis Colon Rectum*; 48 : 1913-6.
6. Karanlik H, Akturk R, Camlica H, Asoglu O. The effect of glyceryl trinitrate ointment on post-hemorrhoidectomy pain and wound healing. Results of a randomised, double-blind, placebo-controlled study. *Dis Colon Rectum* 2009; 52: 280-5.
7. Lunniss P J. The anus and anal canal. In: Williams N S, Bulstrade C J K, Ronan O'Connel P. Eds Bailey and Love's Short Practice of Surgery, London : Edward Arnold; 2008: 1240-70.
8. MacKay D. Hemorrhoids and varicose veins. A review of treatment options. *Altern. Med Rev*. 2001; 6 : 126-40.
9. Kanellos I, Zacharak. J F, Chistoforidis E, Angelopoulos S, Kanellos D, Pramateftakis M G, et al. Usefulness of lateral internal sphincterectomy in reducing post-operative pain after open hemorrhoidectomy. *World J Surg* 2005; 29: 464-8.
10. Rotholtz N A, Bun M, Mauri M V, Bosio R, Peczan C E, Mezzadr N A. Long term assessment of fecal incontinence after lateral internal sphincterectomy. *Tech Colorectal* 2005; 9: 115-18.
11. Hashmi F, Siddiqui F G. Diltiazem 2% versus glyceryl trinitrate cream (0.2%) in the management of chronic anal fissures. *J Coll Physicians Surg Pak* 2009; 19: 750-3.

12. Ganio E, Altomare D F, Gabrielli F, Milito G, Canuti S. Prospective randomised multicenter trial comparing stapled with open hemorrhoidectomy. *Br J Surg* 2001; 88: 669-74.
13. Schonten W R, Van Vroon Hoven T J. Lateral internal sphincterotomy in the treatment of hemorrhoids. A clinical and manometric study. *Dis Colon Rectum* 1986; 29: 869-72.
14. Garcea G, Sutton S, Mansoori S, Lloyd T, Thomas M. Results following conservative lateral sphincterotomy for the treatment of chronic anal fissures. *Colorectal* 2003; 5: 311-4.
15. Galizia G, Lieto E, Imperatore V, Pelosio L, Castellano P. The usefulness of lateral internal sphincterotomy combined with hemorrhoidectomy in the treatment of hemorrhoids: a randomised prospective study. *G Chir* 2000; 21: 27-34.
16. Wasvary H J, Hain J, Moged-Vogel M, Bendick P, Barkel D C, Klein S N. Randomised, prospective, double-blind, placebo-controlled trial of the effects of nitroglycerin ointment on pain after hemorrhoidectomy. *Dis Colon Rectum* 2001; 44: 1069-73.
17. Lund J N, Scholefield J H. A randomised, prospective, double-blind, placebo-controlled trial of glyceryl trinitrate ointment in the treatment of anal fissure. *Lancet* 1997; 349: 11-4.
18. Muhammad O F, Shakir H, Amjad H M, Sohail K M, Zakaria I, Shah A. A role of topical glyceryl trinitrate in the management of anal fissure. *Ann King Edward Med Coll* 2007; 13: 35-8.
19. Jawaid M, Masood Z, Salim M. Topical diltiazem hydrochloride and glyceryl trinitrate in the treatment of chronic anal fissures. *J Coll Physicians and Surgeons Pak* 2009; 19: 614-7.
20. Amoli HA, Notash AY, Shahandashti FJ, Kenari AY, Ashraf H. A randomized, Prospective, Double-blind, Placebo-Controlled Trial of the Effect of Topical Diltiazem on Post-hemorrhoidectomy Pain. *Colorectal Dis* 2009 Nov 14.
21. Chauhan A, Tiwari S, Mishra VK, Bhatia PK. Comparison of internal sphincterotomy with topical Diltiazem for Post-hemorrhoidectomy pain relief: a prospective randomized trial. *J Postgrad Med* 2009; 55: 22-6

ONLINE SUBMISSION OF MANUSCRIPT

It is mandatory to submit the manuscripts at the following website of KJMS. It is quick, convenient, cheap, requirement of HEC and Paperless.

Website: www.kjms.com.pk

The intending writers are expected to first register themselves on the website and follow the instructions on the website. Author agreement can be easily downloaded from our website. A duly signed author agreement must accompany initial submission of the manuscript.